

Topic	IPOL®
Manufacturer	Sanofi Pasteur Prescribing Info (FDA)
Protects Against	Poliovirus
Approved Ages	6 weeks and older
Funding Source	 
Routine Schedule	<p>Children: 4-dose series: 2, 4 & 6-18 months then 4-6 years</p> <p>Adults: <u>Unvaccinated:</u> 0, 1-2 & 6-12 months apart to complete 3-dose series <u>1-2 prior doses:</u> 1-2 doses to complete 3-dose series <u>≥3 prior doses:</u> 1 lifetime booster if at increased risk of exposure</p>
Minimum Intervals	<p>Children:</p> <ul style="list-style-type: none"> • 4 weeks between dose 1 & 2 • 4 weeks between dose 2 & 3 • 6 months between dose 3 & 4 <p>Adults:</p> <ul style="list-style-type: none"> • 4 weeks between dose 1 & 2 • 6 months between dose 2 & 3
Administration	Intramuscular (IM) injection or Subcutaneous (SC) injection
Packaging	Vaccine packaged as a 10-dose multi-dose vial. Use MDV polio until expiration date once punctured.
Storage	Refrigerate between 36°F and 46°F (2°C to 8°C) Protect Vaccine from Light - Do Not Freeze
Recommendations	Polio Vaccine (ACIP)
VFC Letter	Not available
Billing Codes	CPT code for vaccine: 90713 CPT code for administration*: 90460 Medi-Cal Fee-For-Service (FFS) administration: 90713 with modifiers –SK (high-risk) and –SL (VFC) ICD-10-CM code (encounter for immunization): Z23
Comments	Licensed in 1990

Refer to [Vaccine Fact Sheets](https://www.eziz.org/Resources/VaccineFactSheets) (EZIZ.org/Resources/VaccineFactSheets).