

# Respiratory Disease Immunization Recommendations for Children and Adolescents

	July	Aug	Sep	Oct	Nov	Dec	Jan	Feb	Mar	April	May	Jun
<b>COVID-19 (6 months+)</b>	Everyone 6 months and older should have access and the choice to receive COVID-19 vaccine. <a href="#">COVID-19 Vaccination</a> is especially recommended to all children 6-23 months of age and ages 2 years and older with <a href="#">certain risk factors</a> .											
<b>Flu (6 months+)</b>	<b>Vaccinate:</b> <ul style="list-style-type: none"> <li>Children who need 2 doses*</li> <li>Pregnant persons in third trimester</li> <li>Those who may not return in the fall</li> </ul>		<b>Optimal vaccination:</b> <b>September and October</b>		<b>Continue vaccinating as long as flu is circulating, and unexpired vaccine is available.</b>							
<b>RSV (0 -19 months)**</b>				<b>Optimal administration:</b> <b>October 1 – March 31</b>								

Recommended immunization timing     IZ timing for certain situations

\*Children ages 6 months – 8 years who have received less than 2 doses in previous flu seasons, need 2 doses. The first flu vaccine dose should be given as soon as vaccine is available to allow the second dose to be given at least 4 weeks later and ideally by the end of October.

\*\*Infants need RSV immunization if prenatal RSV vaccination status is any of: <14 days before birth, unvaccinated, or unknown. For more details, see [AAP’s Policy Statement on Recommendations for the Prevention of RSV Disease in Infants and Children](#).

This publication was supported by a Centers for Disease Control and Prevention (CDC) grant. Its contents do not necessarily represent the views of CDC.