

Routine Immunization Timing 2026

Suggested schedule to meet recommendations on time.

| |
|---|
| Birth |
| HepB¹ |
| RSV² (age: < 8 months) |

| | |
|--|--|
| 6 months – 18+ years | |
| COVID-19 vaccine(s)⁷ | Flu vaccine, every fall⁸ |

| Age 2 months | Interval from previous dose |
|---|---|
| DTaP (Diphtheria, Tetanus, Pertussis) | |
| Polio (IPV) | |
| HepB³ (age: 1-2 months) | 1-2 months after birth dose |
| Hib⁴ (Hib meningitis) | |
| PCV (Pneumo) | |
| RV⁵ (Rotavirus) | |

| Age 4 months | Interval from previous dose |
|--|--------------------------------------|
| DTaP | 1-2 months |
| Polio (IPV) | 1-2 months |
| HepB³ if 1st dose given at 2 months | 1-2 months |
| Hib | 1-2 months |
| PCV | 1-2 months |
| RV⁵ | 4-10 weeks |

| Age 6 months | Interval from previous dose |
|---|---|
| DTaP | 1-2 months |
| Polio (age: 6-18 months) | 1-14 months |
| HepB³ (age: 6-18 months) | 2-12 months and ≥4 months after 1st dose |
| Hib⁶ | 1-2 months |
| PCV | 1-2 months |
| RV⁵ if RotaTeq used for doses 1 or 2 | 4-10 weeks |

| Age 12 months | Interval from previous dose |
|--|--------------------------------------|
| HepA⁹ (age: 12-23 months) | |
| MMR^{9,10,11} (ages 12-15 months) | |
| Var¹¹ (age: 12-15 months) | |
| Hib (age: 12-15 months) | 2-8 months |
| PCV¹² (age: 12-15 months) | 8 weeks |

| Age 15 months | Interval from previous dose |
|--|--------------------------------------|
| DTaP¹³ (age: 15-18 months) | 6-12 months |

| Age 18 months | Interval from previous dose |
|----------------------------|--------------------------------------|
| HepA | 6-18 months |

| | |
|--------------------------------|---|
| Age 4-6 years | DTaP Polio (IPV) MMR^{10, 11} Varicella¹¹ |
| Age 11-12 years | HPV¹⁴ (2 doses, <u>can</u> <u>start at age 9</u>) MenACWY (MCV4) Tdap |
| Age 16 years | MenACWY (MCV4) MenB¹⁵ |

California Kids
Love them. Immunize them.



This is a suggested schedule. For details, see [AAP's immunization schedule \(PDF\)](#) (AAP.org/immunizationschedule)

1. Monovalent HepB vaccine is recommended within 24 hours of birth for stable infants weighing ≥ 2 kg. For others, see schedule.
2. Recommended from October 1 to March 31. If prenatal RSV vaccination status is any of: <14 days before birth, unvaccinated, or unknown, then infants born during RSV season should receive an RSV immunization as soon as possible after birth, ideally during the birth hospitalization. Infants younger than 8 months born before the RSV season, should receive an RSV immunization before RSV season begins. High-risk children ages 8-19 months should receive nirsevimab before their second RSV season. View [AAP's Policy Statement on Recommendations for the Prevention of RSV Disease in Infants and Children](#) for additional guidance.
3. A dose of HepB vaccine is not necessary at 4 months if doses are given at birth and 2 months but may be included as part of a combination vaccine. The final dose (3RD/4TH) should be given after age 24 wks. and at least 16 wks. after 1st dose.
4. Vaxelis[®], along with PedvaxHIB[®], is preferentially recommended for American Indian & Alaska Native infants for Hib vaccination.
5. Administer first dose at age 6 wks-14 wks. (Max. age: 14 wks., 6 days). Max. age for final dose in the series: 8 months, 0 days. If any dose of RotaTeq[®] (RV5) is given or product is unknown, a total of three RV doses are needed.
6. This 6 month Hib dose is not indicated if PedvaxHIB[®] is used exclusively for the 2 and 4 month infant doses.
7. View pediatric COVID-19 vaccine recommendations on the AAP website and [COVID Vaccine Timing Guide \(PDF\)](#).
8. Two doses given at least 4 weeks apart are recommended for ages 6 months–8 years who are getting flu vaccine for the first time.
9. Refer to CDC guidelines for vaccinating children 6-11 months old prior to international travel. Children 12 months and older should receive 2 doses before international travel.
10. Minimum interval between 1ST and 2ND dose is 4 wks. Two MMR doses should still be given on or after 12 months of age.
11. Varicella vaccine minimum intervals: Ages 1-12 year: 3 months. Ages 13 years and older: 4 weeks. MMRV may be used when both MMR and Varicella vaccines are indicated. For the 1st dose at 12-47 months, MMR and varicella vaccines should typically be given separately unless the parent or caregiver prefers MMRV.
12. Final dose of PCV series should be given at ≥ 12 months of age or after.
13. The 4th dose of DTaP may be administered as early as 12 months, provided at least 6 months have elapsed since the 3RD DTaP dose.
14. HPV vaccine should be given on a 0, 6-12 month schedule for 9-14 year olds (min. interval is 5 months). If patient immunocompromised or initiates series at 15 years or older, use a 3 dose schedule (0, 1-2, 6 months).
15. Shared clinical decision-making is recommended regarding MenB vaccination for healthy people 16–23 years of age. For more details, see [AAP's immunization schedule \(PDF\)](#). Pentavalent MenABCWY vaccine may be used when both MenACWY and MenB are indicated at the same visit. MenB component brands are not interchangeable.